

**PurAmino™ Infant Formula Powder**  
***Dilutions per 1 Scoop***

**Note: Powdered products are not commercially sterile and should not be used for premature or immunocompromised patients, unless clinically required, and then under strict medical supervision of preparation and use.**

Calories Desired,		1 <i>unpacked, level</i> scoop powder added to indication volume of water*		Formula Yield,	
Per fl oz	Per mL	fl oz	mL	fl oz	mL
10	0.34	2.1	63	2.3	67
15	0.51	1.4	41	1.5	44
20	0.68	1.0	30	1.1	33
22	0.74	0.9	27	1.0	30
24	0.81	0.8	24	0.9	28
26	0.88	0.7	22	0.9	26
27	0.91	0.7	21	0.8	25
28	0.95	0.7	20	0.8	24
30	1.01	0.6	19	0.8	22

\*Fluid ounce measure are provided for convenience when larger variance in prepared dilution is acceptable. Fluid ounce measures in above table are rounded to the nearest 0.1 fluid ounce.

When accuracy is essential, water should be measured in mL, and powder should be weighed rather than measured. Contact [mjmedicalaffairs@mjn.com](mailto:mjmedicalaffairs@mjn.com) for those instructions.

One-half fl oz of water can be measured using one tablespoon (1/2 fl oz water = 1 Tbsp water).

Powders mix best when added on *top* of water. Add powder, then close bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One unpacked scoop of PurAmino™ powder (4.5g) provides 23 calories.

100 g of PurAmino™ powder provides 500 Calories. Each gram of powder displaces 0.77 mL water.

One fl oz = 29.57 mL