

## Enfamil NeuroPro™ Infant Formula Powder

### *Dilutions per 1 Scoop*

**Note: Powdered products are not commercially sterile and should not be used for immunocompromised patients, unless clinically required, and then under strict medical supervision of preparation and use.**

Calories Desired, per fl oz	1 Scoop powder added to indication volume of water*, fl oz	Formula Yield, fl oz
20	2.0	2.2
22	1.8	2.0
24	1.6	1.9
26	1.5	1.7
27	1.4	1.7
28	1.4	1.6
30	1.3	1.5

Enfamil NeuroPro™ Infant powder should be measured with **unpacked**, level scoops, using the scoop enclosed in the can.

\*Fluid ounce measure are provided for convenience when larger variance in prepared dilution is acceptable. Fluid ounce measures in above table are rounded to the nearest 0.1 fluid ounce.

One-half fl oz of water can be measured using one tablespoon (1/2 fl oz water = 1 Tbsp water).

When accuracy is essential, water should be measured in mL, and powder should be weighed rather than measured. Contact [mjmedicalaffairs@mjn.com](mailto:mjmedicalaffairs@mjn.com) for those instructions.

Powders mix best when added on *top* of water. Add powder, then close bottle and shake for about 5 seconds. The number of scoops and the amount of water to add can be doubled or tripled to make a larger volume.

One unpacked scoop of Enfamil NeuroPro™ Infant powder (8.8g) provides 45 calories (5.1 kcal/g).

One unpacked scoop of Enfamil NeuroPro™ Infant powder displaces about 0.2 fl oz of water (displacement for Enfamil NeuroPro™ Infant is 0.77 mL/g).

One fl oz = 29.57 mL